arXiv:1207.0416v1 [hep-th] 2 Jul 2012

Some classes of renormalizable tensor models

Joseph Ben Geloun[1, 2, *] and **Etera R. Livine**[3, 1, †]
[1] *Perimeter Institute for Theoretical Physics, 31 Caroline St, Waterloo, ON, Canada*
[2] *International Chair in Mathematical Physics and Applications,*
ICMPA-UNESCO Chair, 072BP50, Cotonou, Rep. of Benin
[3] *Laboratoire de Physique, ENS Lyon, CNRS-UMR 5672, 46 Allée d'Italie, Lyon 69007, France*
(Dated: July 3, 2012)

We identify new families of renormalizable of tensor models from anterior renormalizable tensor models via a mapping capable of reducing or increasing the rank of the theory without having an effect on the renormalizability property. Mainly, the rank 3 tensor model as defined in [arXiv:1201.0176 [hep-th]], the Grosse-Wulkenhaar model in 4D and 2D generate three different classes of renormalizable models.

Pacs numbers: 11.10.Gh, 04.60.-m, 02.10.Ox
Key words: Renormalization, tensor models, matrix models.
pi-qg-285 and ICMPA/MPA/2012/12

I. INTRODUCTION

Recently, a new class of renormalizable models involving tensor fields over $U(1)$ have been highlighted [1–3]. They pertain to the "discrete to continuum" scenario for quantum gravity [4–6]. In their renormalization procedure, the multiscale analysis [7] led to a new kind power counting theorem and locality principle (using the correct extension of $1/N$ expansion for higher rank theories [8, 9]) shedding more light on anterior results in renormalization of matrix models [10–12]. At the perturbative level, Feynman graphs of these models are generated by vertices spanned by nonlocal interactions representing simplices glued by higher rank extension of ribbon propagators.

Hence, in a broader sense, the models cited above belong to the class of models endowed with nonlocal interactions. Field arguments in the interaction term might be paired in many possible ways. Specific forms have to be physically motivated. For instance, in noncommutative field theory induced by noncommutative Moyal field algebra, four fields interact in a region (a parallelogram) the area of which is the Planck length square [10]. The recent tensor models in [1, 2] possess interactions of the form of 3 and 4-simplex which generate, through a path integral formalism, simplicial pseudo-manifolds in 3D and 4D. They should be regarded as the higher rank extension of the celebrated matrix models [13] or rank 3 tensor models [14] with a bonus: the genuine feature to be renormalizable.

In nonlocal field theory, it could happen that the interaction is of a particular form that can be called "partially cyclic", namely, in the interaction pairing, a tensor field only share (at least two) arguments with at most two other fields (a "totally cyclic" or simply "cyclic" interaction would be an interaction having this property valid for all fields which define it). Roughly, consider some tensors T^i and an interaction $I = \sum_{[J],[K],\dots} T^1_{\dots,[J],\dots} T^2_{\dots,[J],[K],\dots} T^3_{\dots,[K],\dots} \cdots$, where $[J]$ and $[K]$ are block indices or arguments and a simpler interaction of the form $\tilde{I} = \sum_{j,k,\dots} T^1_{\dots,j,\dots} T^2_{\dots,j,k,\dots} T^3_{\dots,k,\dots}$ .... It is natural to ask: how starting from I, one ends up with $\tilde{I}$ and what is the main feature of the reduced model described by $\tilde{I}$? A quantum field theory being not only defined by interactions what implications has such a reduction on the dynamics? These questions might be very intricate and, more to the point, even more complex if one would like to preserve some nice properties (for instance renormalizability or symmetry aspects) of the initial theory.

In this paper, we show that, at least three particular classes of tensor models equipped with cyclic interactions can be projected back to reduced rank models and, reciprocally, any tensor model of this kind can be extended to a higher rank tensor model with a cyclic interaction in a sector. During the process, we are able to identify new classes of renormalizable models (at all orders). If the projection and extension of these models can be somehow understood either by dividing or by multiplying the number of indices, what we actually show is that this mapping preserves the renormalizability of the starting model. The proof is established starting from the rank 3 tensor model of [2], the Grosse-Wulkenhaar (GW) model in 2D [12] and 4D [11], and, from these, the universal feature of our formalism can be easily inferred. The interpretation of the reduced or extended models in terms of the initial model framework will be discussed elsewhere. We stress also the fact that the projection-extension mechanism can be used to reduce a piece

---

[*] jbengeloun@perimeterinstitute.ca
[†] etera.livine@ens-lyon.fr

of the rank 4 model of [1] but cannot reduce the entire interaction to a matrix interaction. Hence, there are actually renormalizable models which are not partially cyclic.

The plan of this paper is the following: In the next section, we discuss slightly in detail, as a test of the above ideas, the rank 3 renormalizable model of [3] and its reduction to a new matrix model. The reasons why the reduced model is renormalizable are determined. Section III is devoted to the identification of new families of tensor models issued from GW models in 4D and 2D which should not be considered as belonging to the same family according to our mapping. We infer the existence of three different families of renormalizable models (rank 3 tensor, GW 4D and GW 2D models) having three different renormalizable matrix models as roots. Interestingly, the rank 3 tensor model "lies in between" (in some sense) the GW 4D and GW 2D models.

II. MATRIX ANALOGUE OF A RANK THREE TENSOR MODEL

Consider the model as defined in [2] henceforth called $T3$ model. Fields $\varphi : U(1)^3 \to \mathbb{C}$ can be equivalently described after Fourier mode decomposition as tensors $\varphi_{[p]}$, $[p] = (p_1, p_2, p_3)$, $p_i \in \mathbb{Z}$. We will restrict the discussion for positive mode fields, i.e. tensor fields satisfying the symmetry

$$\varphi_{p_1,p_2,p_3} = \varphi_{-p_1,p_2,p_3} = \varphi_{p_1,-p_2,p_3} = \varphi_{p_1,p_2,-p_3} \,. \tag{1}$$

Thus, we will consider only fields such that $p_i \in \mathbb{N}$, namely fields can be regarded as living in $(U(1)/\mathbb{Z}_2)^3$. Note such a restriction is made by sake of simplicity. Hence, such a prescription will have no consequence on the subsequent analysis (this point will be emphasized later on).

Using now these tensor components such that $\varphi_{p_1,p_2,p_3} \in \mathbb{C}$, $p_i \in \mathbb{N}$, the $T3$ model possesses a kinetic term given by

$$S^{\rm kin}_{T3} = \sum_{p_j} \bar\varphi_{p_1,p_2,p_3} \Big(\sum_{s=1}^{3} p_s + \mu \Big) \varphi_{p_1,p_2,p_3} \,, \tag{2}$$

with some mass μ, associated with a propagator $\widehat{C}([p_i];[\tilde p_i]) = \prod_{i=1}^{3} \delta_{p_i,\tilde p_i} / \big(\sum_{s=1}^{3} p_s + \mu \big)$ and a φ^4 interaction of the form

$$S^{\rm int}_{T3} = \sum_{p_j} \varphi_{p_1,p_2,p_3}\, \bar\varphi_{p_{1'},p_2,p_3}\, \varphi_{p_{1'},p_{2'},p_{3'}}\, \bar\varphi_{p_1,p_{2'},p_{3'}} \,. \tag{3}$$

Propagator and vertex are given in Fig.1. Note that the $T3$ model as introduced in [2] involves more interactions.



FIG. 1. Propagator and vertex of the type φ^4 in rank 3 tensor model.

Nevertheless, an important feature of this model is that the flow of each interaction does not generate any other coupling than itself. The Renormalization Group (RG) equations at the first order of perturbation theory have displayed this property [2] however a simple proof of this claim will be given below. Thus, given this feature, we will restrict to the renormalizable model with the unique interaction (3).

A striking feature of the interaction (3) is that it can be mapped to a pure matrix interaction using any bijection $\tilde\sigma : \mathbb{N}^2 \to \mathbb{N}$. Indeed, consider the following field redefinition[1]:

$$\varphi_{p_1,p_2,p_3} \;\mapsto\; \phi_{p,n} \,, \qquad p_1 = p \,, \quad \tilde\sigma(p_2,p_3) = n \tag{4}$$

---

[1] We reduce the (p_2, p_3)-momentum sector according to the fact that the interaction is cyclic with respect to this couple of indices.

to which, given $\tilde\sigma(p,q)=n$ and its inverse noted as $\tilde\sigma_1^{-1}(n)=p$ and $\tilde\sigma_2^{-1}(n)=q$, corresponds the following transformed actions

$$S^{\rm kin}_{T3} = \sum_{p,n} \bar\phi_{p,n}\Big(p+\tilde\sigma_1^{-1}(n)+\tilde\sigma_2^{-1}(n)+\mu\Big)\phi_{p,n}\,, \tag{5}$$

$$S^{\rm int}_{T3} = \sum_{p_1,p_2,n_1,n_2} \phi_{p_1,n_1}\,\bar\phi_{p_2,n_1}\,\phi_{p_2,n_2}\,\bar\phi_{p_1,n_2}\,. \tag{6}$$



FIG. 2. Propagator and vertex in the reduced matrix theory.

The interaction (6) is clearly of the matrix kind and cannot generate by RG flow any other coupling than itself. It can be represented in the ordinary form of Fig.2. Since this is still the same model, the renormalizability of the model (5)-(6) is guaranteed. However, it can be instructive to scrutinize again the behavior of the propagator in this reshuffled basis. To this end, using Schwinger's kernel and $N_n=\tilde\sigma_1^{-1}(n)+\tilde\sigma_2^{-1}(n)$, we write the slice decomposition of the propagator kernel in the standard way of [7] as

$$\begin{aligned}
C(p,n) &= \frac{1}{p+N_n+\mu} = \int_0^\infty d\alpha\, e^{-\alpha(p+N_n+\mu)}\,,\\
C_i(p,n) &= \int_{M^{-i}}^{M^{-i+1}} d\alpha\, e^{-\alpha(p+N_n+\mu)} \leq K M^{-i} e^{-\delta M^{-i}(p+N_n+\mu)}\,,\\
C_0(p,n) &= \int_1^\infty d\alpha\, e^{-\alpha(p+N_n+\mu)} \leq K e^{-\delta(p+N_n+\mu)}\,,
\end{aligned} \tag{7}$$

for some large constant $M\in\mathbb{N}$, $K\geq 0$ and $\delta\geq 0$. One has $C=\sum_{i=0}^\infty C_i$. Hence, a slice $i\gg 1$ probe high momenta. A UV cut-off can be introduced such that $C^\Lambda=\sum_{i=0}^\Lambda C_i$ and the UV limit is obtained by taking $\Lambda\to\infty$.

In particular, the standard choice σ for $\tilde\sigma$ is defined by ordering the pairs of integers (p,q) in $\mathbb{N}^2$ along the diagonal (at constant $p+q$) and numbering them from bottom to top, as illustrated in Fig.3.



FIG. 3. On the left, the bijection σ labeling all pairs of integers and mapping $\mathbb{N}^2$ onto $\mathbb{N}$, and on the right its generalization to a bijective map from $\mathbb{Z}^2$ onto $\mathbb{N}$.

Explicitly, this map reads:

$$\sigma(p,q) = \frac{1}{2}(p+q)(p+q+1)+q \tag{8}$$

and its inverse can be characterized as

$$\sigma^{-1}(n) = (p,q)\in\mathbb{N}\,,\quad N=p+q\,,\quad N(N+1)\leq 2n\leq N(N+3)\,,\quad q=2n-N(N+1)\,,\quad p=N-q\,, \tag{9}$$

where the inequality uniquely determines $N=N_n\in\mathbb{N}$. We denote $\sigma_1^{-1}(n)=p$ and $\sigma_2^{-1}(n)=q$. It can be easily shown that the following (optimal) bounds holds:

$$\frac{\sqrt{9+8n}-3}{2}\leq N\leq\frac{\sqrt{1+8n}-1}{2}\,,$$

$$\frac{(\sqrt{9+8n}-3)^2}{8} \leq \frac{N^2}{2} \leq p^2+q^2 \leq \Big[(p+q)^2 = N^2\Big] \leq \frac{(\sqrt{1+8n}-1)^2}{4} \leq 2n\,. \tag{10}$$

Hence, for large n,

$$\sqrt{2n}-\frac{3}{2} \leq \sqrt{2n}\Big[\sqrt{\frac{9}{8n}+1}-\frac{3}{\sqrt{8n}}\Big] \leq N \leq \frac{\sqrt{1+8n}-1}{2} \leq \sqrt{2n}\,, \tag{11}$$

the approximation $N \sim \sqrt{2n}$ is correct. We could of course introduce another choice of bijection between $\mathbb{N}^2$ and $\mathbb{N}$, but the map σ is, in some sense, the most compact choice for which we have a natural simple estimate of $N = p+q$ in terms of $n = \sigma(p,q)$. For a more general map, we would reshuffle the labeling of points $(p,q) \in \mathbb{N}^2$ by the integer n, which would lead to a more random behavior of N that would wildly fluctuate away from $\sqrt{2n}$.

Let us point out that we can easily adapt our analysis to $\mathbb{Z}^2$ and introduce a generalized map $\sigma : \mathbb{Z}^2 \to \mathbb{N}$ as shown in Fig.3, labeling points $(p,q) \in \mathbb{Z}^2$ along the "circles" of constant $N = |p|+|q|$. This would lead to a similar behavior of N scaling proportionality to $\sqrt{n}$. This also sustains the fact that we could have let all field modes $p_i \in \mathbb{Z}$ (without assuming any symmetry (1)) and still the analysis will be valid.

Coming back to the propagator and using the fact that there exists a constant k such that $k\sqrt{n} \leq N$ (for instance N is larger than $\sqrt{n/2}$ as soon as $n \geq 2$), we get the following bound on the sliced propagator:

$$C_i(p,n) \leq KM^{-i}e^{-\delta' M^{-i}(p+\sqrt{n}+m)}\,. \tag{12}$$

We can now infer that there is a new model that we will call $\sigma T3$ with the same interaction $\widetilde{S}^{\rm int}_{T3} = S^{\rm int}_{T3}$ and the following kinetic term

$$\widetilde{S}^{\rm kin}_{T3} = \sum_{p,n} \bar{\phi}_{p,n}\Big(p+\sqrt{n}+m\Big)\phi_{p,n}\,, \tag{13}$$

such that the action $\widetilde{S} = \widetilde{S}^{\rm kin}_{T3} + (\lambda/4)\widetilde{S}^{\rm int}_{T3}$ is renormalizable at all orders of perturbation theory.

Due to the particular form of the propagator and vertex (a product of delta functions in the momentum basis), the divergences occurring in graph amplitudes in the above models come only from closed loops or internal faces (in the momentum basis). The amplitude of these faces can be recast as sums of the form:

$$\sum_p e^{-\delta M^{-i}p} = \delta' M^i + O(M^{-i})\,, \qquad \sum_n e^{-\delta M^{-i}\sqrt{n}} = \delta' M^{2i} + O(M^i)\,, \tag{14}$$

where δ and δ' are some constants, and we consider large scale $i \gg 1$.

We can now understand the main feature introduced the field redefinition. After monoscale expansion[2] of the Feynman amplitude $A_{\mathcal{G}} = \sum_i A_{\mathcal{G};i}$ of a given graph $\mathcal{G}$, the $T3$ model generates a crude power-counting yielding a divergence degree $\omega_d(\mathcal{G}) = -L(\mathcal{G}) + F_{\rm int}(\mathcal{G})$, where $L(\mathcal{G})$ is the number of lines and $F_{\rm int}(\mathcal{G})$ the number of closed strands or internal faces of the graph $\mathcal{G}$ [2]. We have $A_{\mathcal{G};i} = KM^{i\omega_d(\mathcal{G})}$. After mapping $T3 \to \sigma T3$, one realizes that, from (12), each line l at scale i_l provides a convergent factor of M^{-i_l} yielding, roughly, a convergent factor of $M^{-iL(\mathcal{G})}$, for a monoscale expansion. This is exactly similar to the situation of the $T3$ model. In contrast, the number of internal faces has been drastically reduced by merging the strands after the rank reduction. There are two types of internal faces: those indexed by p_f in $\mathcal{F}^-_{\rm int}(\mathcal{G})$ (of cardinal $F^-_{\rm int}(\mathcal{G})$) which yield a divergent factor M^i (14); and internal faces labeled by $\sqrt{n_f}$ in $\mathcal{F}^+_{\rm int}(\mathcal{G})$ (of cardinal $F^+_{\rm int}(\mathcal{G})$) yielding a greater divergent factor of M^{2i} from (14). The latter faces thereby compensate the missing faces after the tensor to matrix reduction and, from that, one recovers the balanced power-counting theorem of the initial model. In symbol, we write:

$$\omega_d(\mathcal{G}) = -L(\mathcal{G}) + F^-_{\rm int}(\mathcal{G}) + 2F^+_{\rm int}(\mathcal{G}) = -L(\mathcal{G}) + F_{\rm int}(\mathcal{G}) + F^+_{\rm int}(\mathcal{G}) \leq -L(\mathcal{G}) + 2F_{\rm int}(\mathcal{G})\,. \tag{15}$$

The bound by $-L(\mathcal{G}) + 2F_{\rm int}(\mathcal{G})$ is, in fact, optimal since there are some configurations where $F_{\rm int}(\mathcal{G}) = F^+_{\rm int}(\mathcal{G})$. Henceforth, we will use the divergence degree $\omega_d(\mathcal{G}) = -L(\mathcal{G}) + 2F_{\rm int}(\mathcal{G})$. Note that the above contribution $2F^+_{\rm int}(\mathcal{G})$ can be inferred either by straightforward calculations or understood from the fact that faces in $\mathcal{F}^+_{\rm int}$ have to be simply regarded as double faces.

---

[2] A detailed multiscale expansion can be performed as well and the following will be valid.

It is worth mentioning that the detailed proof of renormalizability in the present setting would be an interesting task even though somehow beyond the scoop of this paper. Nevertheless, using the same techniques used in [1] and [2], we can recast (15) in terms of topological numbers as

$$\omega_d(\mathcal{G}) = -V_2 - \frac{1}{2}\big[N_{\text{ext}} - 4\big] - 4g_{\widetilde{\mathcal{G}}} - 2(C_{\partial\widetilde{\mathcal{G}}} - 1)\,, \tag{16}$$

where V_2 is the number of mass renormalization vertices, $g_{\widetilde{\mathcal{G}}}$ is the genus of the closed colored[3] graph [15, 16] $\widetilde{\mathcal{G}}$ extending the initial graph $\mathcal{G}$, $C_{\partial\widetilde{\mathcal{G}}}$ is the number of connected components of $\partial\widetilde{\mathcal{G}}$ the boundary graph associated with $\widetilde{\mathcal{G}}$ (or collection of all "pinched" external faces; for a complete definition of $\widetilde{\mathcal{G}}$ and $\partial\widetilde{\mathcal{G}}$, we shall refer the reader to [1] or [2]) and N_{ext} is the number of external legs of $\mathcal{G}$. From (16), one realizes that the marginal graph category or log-divergent graphs ($\omega_d(\mathcal{G}) = 0$) with $N_{\text{ext}} = 4$ are planar ones with $C_{\partial\widetilde{\mathcal{G}}} = 1$ and $V_2 = 0$. They should correspond to a vertex renormalization. For $N_{\text{ext}} = 2$, there are two divergent categories: linearly divergent graphs ($\omega_d(\mathcal{G}) = 1$) are those with, $C_{\partial\widetilde{\mathcal{G}}} = 1$ and $V_2 = 0$ and log-divergent contributions with $C_{\partial\widetilde{\mathcal{G}}} = 1$ and $V_2 = 1$ which are subleading. These correspond to a mass renormalization and wave function renormalization for $\omega_d(\mathcal{G}) = 1$. The corresponding subtractions and proof that the divergent parts of N_{ext}-point functions are of the form of the initial Lagrangian terms can be performed in the way of [2]. Note that it is important that we have chosen a kinetic term of form $p^\alpha + q^\beta$ for obtaining the correct wave function counterterms. We remark also that this power-counting is totally similar of the one of the GW model in 4D for which the renormalization procedure identifies as relevant graphs only planar graphs with one broken external face with at most four external legs [11].

There is a now a clear effect induced by the projection map. In the renormalization procedure of $T3$ [2], the dominant contributions, at given $N_{\text{ext}} = 4, 2$, were specific graphs called "melonic"[4] [9, 17, 18] with melonic boundary graph $\partial\widetilde{\mathcal{G}}$ having a unique connected component. However, a look on the divergence degree $\omega_d(\mathcal{G})$ (16), one realizes the fact that dominant contributions should be melonic has been washed away and gets replaced just by a planarity condition. This can be indeed expected since the new model is a matrix theory which can only be sensitive to an ordinary planarity condition. The above discrepancy can be also explained by the fact that we have performed an approximation $F^+_{\text{int}} \simeq F_{\text{int}}$ and, from that, have used $\omega_d(\mathcal{G}) = -L(\mathcal{G}) + 2F_{\text{int}}(\mathcal{G})$. Nevertheless, for melonic graphs F^+_{int} should be the one which contributes so that one would rather have a power counting much more restrictive and of the form $\widetilde{\omega}_d(\mathcal{G}) = -L(\mathcal{G}) + 2F^+_{\text{int}}(\mathcal{G})$ so that many planar graphs would be cast away from being convergent with respect to this power counting.

III. NEW CLASSES OF RENORMALIZABLE MODELS FROM PREVIOUS MODELS

In this section, we use the above mechanism to reveal the existence of new renormalizable models issued from well-known renormalizable higher rank tensor models. Even though the renormalizability of these new reduced model can be regarded as granted by the one-to-one map, the full-fledged proof of renormalizability in the new basis should be considered as an investigation on its own. We will provide the underpinning arguments of why the renormalization should hold.

A. Classes of GW models

The GW model is the first renormalizable model pertaining to both matrix models and noncommutative geometry [11][12]. This model proves to be renormalizable at all orders by curing a previous undesirable effect called UV/IR mixing affecting renormalization procedure in noncommutative spaces [10]. The UV/IR mixing is simply removed by adding an harmonic term of the form $\Omega\tilde{x}^2$, where Ω is an harmonic frequency, $\tilde{x} = 2(\Theta^{-1})_{\mu\nu}x^\mu$, with the noncommutative structure given by $[x^\mu, x^\nu] = \Theta^{\mu\nu}$, $x^\mu \in \mathbb{R}^4$.

We will restrict the study to complex fields and will place ourselves at the self-dual point, $\Omega = 1$ for which, in the continuum, the kinetic term of the GW model is $(-\Delta + \tilde{x}^2 + \mu)$, μ being some IR mass regulator, so that the model

---

[3] Colored graphs in the sense of Gurau [15, 16] are graphs defined by the path integral of the following action for rank 3 fields:

$$S_{\text{color}} = \sum_{p_i}\sum_{a=0}^{3} \bar{\varphi}^a_{p_1,p_2,p_3}\varphi^a_{p_1,p_2,p_3} + \sum_{p_i}\Big[\bar{\varphi}^0_{p_1,p_2,p_3}\bar{\varphi}^1_{p_3,p_4,p_5}\bar{\varphi}^2_{p_5,p_2,p_6}\bar{\varphi}^3_{p_6,p_4,p_1} + \varphi^0_{p_1,p_2,p_3}\varphi^1_{p_3,p_4,p_5}\varphi^2_{p_5,p_2,p_6}\varphi^3_{p_6,p_4,p_1}\Big]\,, \tag{17}$$

where $a = 0, 1, 2, 3$ is referred to as the color index. Such a model admits an extension for arbitrary rank fields.

[4] Roughly speaking, a "melonic" colored tensor graph has all genera $g_{\widetilde{J}} = 0$ for all of its pinched (or closed) jackets $\widetilde{J}$. A jacket is a subribbon graph (inside the tensor graph) defined by a cycle of colors of the initial colored graph. In a rank 3 model, $\widetilde{\mathcal{G}}$ is a colored graph with 4 colors 0,1,2,3. There are three different jackets defined by the cycles (0123), (0132) (0213). Note that "melonicity" is a much more stringent condition than the planarity condition on the graph [9, 17, 18]. A melonic graph in dimension D is topologically homeomorphic to the S^D sphere, the converse is not always true.

becomes dual in momenta and positions. In 4D, the complex GW model is given by the action (Euclidean signature) [11]

$$S_{GW;4D} = \int d^4x \left\{ \frac{1}{2}\bar\phi(x)(-(\partial_\nu)^2 + \tilde{x}^2 + \mu)\phi(x) + \frac{\lambda}{4}\bar\phi(x)\star\phi(x)\star\bar\phi(x)\star\phi(x) \right\}, \tag{18}$$

where $\star$ denotes the Moyal star product. Since each field can be viewed as a rank 4 complex tensor $\phi_{\vec p,\vec q}$ where $\vec p = (p_1, p_2)$ and $\vec q = (q_1, q_2) \in \mathbb{N}^2$, the same GW action translated in the matrix basis reads [11]

$$S_{GW;4D} = \frac{1}{2}\sum_{\vec p,\vec q\in\mathbb{N}^2} \bar\phi_{\vec p,\vec q}\Big[p_1+p_2+q_1+q_2+\mu\Big]\phi_{\vec q,\vec p} + \frac{\lambda}{4}\sum_{\vec m,\vec n,\vec p,\vec q\in\mathbb{N}^2} \bar\phi_{\vec m,\vec n}\,\phi_{\vec n,\vec p}\,\bar\phi_{\vec p,\vec q}\,\phi_{\vec q,\vec m}\,, \tag{19}$$

with the last interaction clearly of the cyclic form. Call this the GW_{4D} model.

The previous discussion leads us to consider a new rank 3 model, called σGW_{4D}, described by the following action[5]:

$$\widetilde{S}_{GW;4D} = \frac{1}{2}\sum_{p\in N,\vec q\in\mathbb{N}^2} \bar\phi_{p,\vec q}\Big[\sqrt{p}+q_1+q_2+\mu\Big]\phi_{\vec q,p} + \frac{\lambda}{4}\sum_{m,p\in\mathbb{N};\ s\vec n,\vec q\in N^2} \bar\phi_{m,\vec n}\,\phi_{\vec n,p}\,\bar\phi_{p,\vec q}\,\phi_{\vec q,m}\,. \tag{20}$$

This model is renormalizable at all orders. The proof of this statement goes under the same arguments as previously. Indeed, using the mapping σ in $\vec p$-sector, i.e. $\sigma(p_1, p_2) = p \in \mathbb{N}$, we redefine the field $\phi_{\vec p,\vec q}$ merely as $\phi_{p,\vec q}$. One gets from (19) the reduced rank three action:

$$S_{GW;4D} = S'_{GW;4D} = \frac{1}{2}\sum_{p,\vec q\in\mathbb{N}} \bar\phi_{p,\vec q}\Big[\sigma_1^{-1}(p)+\sigma_2^{-1}(p)+q_1+q_2+\mu\Big]\phi_{\vec q,p} + \frac{\lambda}{4}\sum_{m,p\in\mathbb{N};\ \vec n,\vec q\in N^2} \bar\phi_{m,\vec n}\,\phi_{\vec n,p}\,\bar\phi_{p,\vec q}\,\phi_{\vec q,m}\,. \tag{21}$$

Note that both (20) and (21) share the same interaction. Moreover, in the slice i, the propagator kernel of both models can be bounded by

$$\tilde{C}_i(p,n) \le KM^{-i}e^{-\delta' M^{-i}(\sqrt{p}+q_1+q_2+\mu)}\,. \tag{22}$$

It becomes immediate that σGW_{4D} is renormalizable. A closer inspection on (22) shows that even though the number of faces of the reduced theory certainly decreases in p-sector, the amplitude of each face becomes twice greater in that sector ensuring again the renormalizability.

Applying another reduction in the remaining sector $\vec q$, we get finally a matrix model that we call $\sigma^2 GW_{4D}$ given by

$$\widetilde{\widetilde{S}}_{GW;4D} = \frac{1}{2}\sum_{p,q\in\mathbb{N}} \bar\phi_{p,q}\Big[\sqrt{p}+\sqrt{q}+\mu\Big]\phi_{p,q} + \frac{\lambda}{4}\sum_{m,n,p,q\in\mathbb{N}} \bar\phi_{m,n}\,\phi_{n,p}\,\bar\phi_{p,q}\,\phi_{q,m}\,, \tag{23}$$

the propagator of which scales like the one of the following action

$$S_{GW;4D} = S''_{GW;4D} = \frac{1}{2}\sum_{p,q\in\mathbb{N}} \bar\phi_{p,q}\Big[\sigma_1^{-1}(p)+\sigma_2^{-1}(p)+\sigma_1^{-1}(q)+\sigma_2^{-1}(q)+\mu\Big]\phi_{p,q} + \frac{\lambda}{4}\sum_{m,n,p,q\in\mathbb{N}} \bar\phi_{m,n}\,\phi_{n,p}\,\bar\phi_{p,q}\,\phi_{q,m} \tag{24}$$

and therefore $\sigma^2 GW_{4D}$ is renormalizable at all orders.

It is noteworthy that $\widetilde{\widetilde{S}}_{GW;4D}$ does not correspond to the action of the GW model in 2D, denoted GW_{2D}, which is, in fact, super-renormalizable. Indeed, the action of the latter model is given by [12]:

$$S_{GW;2D} = \frac{1}{2}\sum_{p,q\in\mathbb{N}} \bar\phi_{p,q}\Big[p+q+\mu\Big]\phi_{p,q} + \frac{\lambda}{4}\sum_{m,n,p,q\in\mathbb{N}} \bar\phi_{m,n}\,\phi_{n,p}\,\bar\phi_{p,q}\,\phi_{q,m}\,. \tag{25}$$

Reshuffling now this action by using the inverse of σ^{-1} in one sector, say q without loss of generality, $\sigma^{-1}(q) = \vec q$, we get a rank 3 (super-)renormalizable GW model:

$$S_{GW;2D} = S'_{GW;2D} = \frac{1}{2}\sum_{p\in\mathbb{N},\vec q\in\mathbb{N}^2} \bar\phi_{p,\vec q}\Big[p+\sigma(\vec q)+\mu\Big]\phi_{\vec q,p} + \frac{\lambda}{4}\sum_{m,p\in\mathbb{N};\ \vec n,\vec q\in\mathbb{N}^2} \bar\phi_{m,\vec n}\,\phi_{\vec n,p}\,\bar\phi_{p,\vec q}\,\phi_{\vec q,m} \tag{26}$$

---

[5] Note that, in the new action, the mapping σ can be applied as well on the second couple of integers $\vec q$, without loss of generality. Hence, combinatorially, we have two such σGW_{4D} models.

which can be related to the new $\sigma^{-1}GW_{2D}$ model defined as

$$\widetilde{S}_{GW;2D} = \frac{1}{2}\sum_{p\in\mathbb{N},\vec{q}\in\mathbb{N}^2} \bar{\phi}_{p,\vec{q}}\Big[p + q_1^2 + q_2^2 + \mu\Big]\phi_{p,\vec{q}} + \frac{\lambda}{4}\sum_{m,p\in\mathbb{N};\ \vec{n},\vec{q}\in\mathbb{N}^2} \bar{\phi}_{m,\vec{n}}\,\phi_{\vec{n},p}\,\bar{\phi}_{p,\vec{q}}\,\phi_{\vec{q},m}\,. \tag{27}$$

One can check that, using (10), that the propagators (26) and (27) have same behavior in a slice. In this specific instance, the fact that the number of faces in q-sector is increasing is merely compensated by the fact that each face will be associated with a less divergent factor of $M^{-i/2}$.

Applying now σ^{-1} in the p-sector from (26), we can infer that the following rank 4 GW model which will be referred to as $\sigma^{-2}GW_{2D}$ is (super-)renormalizable as well:

$$\widetilde{\widetilde{S}}_{GW;2D} = \frac{1}{2}\sum_{\vec{p},\vec{q}\in\mathbb{N}^2} \bar{\phi}_{\vec{p},\vec{q}}\Big[p_1^2 + p_2^2 + q_1^2 + q_2^2 + \mu\Big]\phi_{\vec{p},\vec{q}} + \frac{\lambda}{4}\sum_{\vec{m},\vec{n},\vec{p},\vec{q}\in\mathbb{N}^2} \bar{\phi}_{\vec{m},\vec{n}}\,\phi_{\vec{n},\vec{p}}\,\bar{\phi}_{\vec{p},\vec{q}}\,\phi_{\vec{q},\vec{m}}\,. \tag{28}$$

It can be asked the continuum models underlying (20), (23), (27) and (28) and their relation to noncommutative geometry. At this point, an answer to that question is not clear. One can investigate the particular forms of the propagators which might lead to other interesting kinetic terms extending the ordinary (p,x)-duality which has led to the control of UV/IR mixing. Furthermore, if these actions do not find a clear interpretation in Moyal noncommutative field theory, they might be interpreted in some new context of nonlocal field theories. Indeed, in [1], a rank 4 tensor model extending the above $T3$ tensor model has been proved to be renormalizable at all order of perturbation theory. Fields $\varphi : U(1)^4 \to \mathbb{C}$ can be viewed as rank four tensors $\varphi_{p_1,p_2,p_3,p_4}$, $p_i \in \mathbb{Z}$. The kinetic part of this model is given by closely related to the kinetic part of (28). However, the interactions of these two models are different. In [1], one type of interaction is cyclic (hence can be recast in a matrix form) and another cannot be recast in terms of matrix trace. This makes the rank 4 non trivial with this respect. In summary, the type of renormalizable models introduced here might be significant in tensor field theory framework [6].

B. Families of renormalizable tensor models

By iterating the procedure, we can generate three different families of renormalizable models related either to the GW models or to the $T3$ model. We establish that, for the three models,

$$\begin{aligned}
&T3 \text{ Class}: && \cdots \to \sigma^{-n}T3 \to \cdots \to \sigma^{-1}T3 \to T3 \to \sigma T3\,;\\
&GW_{4D} \text{ Class}: && \cdots \to \sigma^{-n}GW_{4D} \to \cdots \to \sigma^{-2}GW_{4D} \to \sigma^{-1}GW_{4D} \to GW_{4D} \to \sigma GW_{4D} \to \sigma^2 GW_{4D}\,;\\
&GW_{2D} \text{ Class}: && \cdots \to \sigma^{-n}GW_{2D} \to \cdots \to \sigma^{-2}GW_{2D} \to \sigma^{-1}GW_{2D} \to GW_{2D}\,.
\end{aligned} \tag{29}$$

Note that each arrow might lead to different theories according to the choice of indices on which the reduction or extension are performed. For instance, $T3 \to \sigma T3$ leads to a unique model whereas $GW_{4D} \to \sigma GW_{4D} \to \sigma^2 GW_{4D}$ leads to two models and $\sigma^{-2}GW_{2D} \leftarrow \sigma^{-1}GW_{2D} \leftarrow GW_{2D}$ leads as well to two models. A way to classify all these models might be to consider as belonging to the same family or class those having a common and initial matrix model.

Being interested in the change of the propagator for these different theories, we have the following table:

Matrix	Rank 3	Rank 4
$\sigma^2 GW_{4D} : \sqrt{p} + \sqrt{q}$	$\sigma GW_{4D} : \sqrt{p} + q_1 + q_2$	$GW_{4D} : p_1 + p_2 + q_1 + q_2$
$\sigma T3 : p + \sqrt{q}$	$T3 : p + q_1 + q_2$	$\sigma^{-1}T3 : p_1^2 + p_2^2 + q_1 + q_2$
$GW_{2D} : p + q$	$\sigma^{-1}GW_{2D} : p + q_1^2 + q_2^2$	$\sigma^{-2}GW_{2D} : p_1^2 + p_2^2 + q_1^2 + q_2^2$

IV. OUTLOOK

The present work affords a link between different nonlocal renormalizable theories using tensor fields of rank ≥ 2 appearing in different field theory contexts (from noncommutative field theory to tensor models). It provides also a machinery in order to generate classes of models sharing the significant property to be renormalizable.

Remarkably, the renormalizability of the $T3$ model can be understood from the fact that, its multi-scale analysis can be performed in the reduced matrix model $\sigma T3$ for which the propagator in a given slice i contains $\sqrt{q}$ which allows to have a similar power counting of $\sigma^2 GW_{4D}$ (of which the renormalizability holds from the one of GW_{4D}). Furthermore, since GW_{2D} is super-renormalizable [12], by introducing a set of propagators $p + q^{\frac{1}{2}+\epsilon}$, $\epsilon \in [0, 1/2]$, we find a continuum of theories interpolating between GW_{2D} for $\epsilon = 1/2$ and $T3$ for $\epsilon = 0$, all with the property from being super-renormalizable to being just-renormalizable. Pursuing the propagator interpolation $p^{\frac{1}{2}+\epsilon'} + q^{\frac{1}{2}}$, $\epsilon' \in [0, 1/2]$, we find another continuum of theories being all just-renormalizable leaving from $T3$ at $\epsilon' = 1/2$ to GW_{4D} at $\epsilon' = 0$. The $T3$ model is, with this respect, "critical". One may also ask about the UV behavior of these classes of models. It can be shown that, all the models such that $p^{\frac{1}{2}+\epsilon} + q^{\frac{1}{2}}$, for $\epsilon \in]0, 1/2]$, are asymptotically free in the UV (see the appendix for a proof of this claim) and, at the end-point $\epsilon = 0$, the model becomes safe which corresponds, of course, of the well-known asymptotic safeness of the GW model in 4D [11][10]. In this sense, GW_{4D} becomes "critical".

Finally, it would be interesting to provide a space-time representation to all these matrix/tensor theories, the same way that the GW model is defined by the action (18) as a field theory living on non-commutative $\mathbb{R}^4$. Since terms in $\sqrt{p}$ in the propagator seem slightly awkward to translate as differential operators (introducing another type of nonlocality), it seems more natural to work with the theories with linear propagator, for instance $p_1 + p_2 + q_1 + q_2$ for GW_{4D} and $p + q_1 + q_2$ for $T3$. Considering $T3$, one can write it naturally as a group field theory [4] on $U(1)^3$ with the kinetic term given by the sum of the derivative with respect to each coordinate. Written as such, we lose a priori the relation with non-commutative field theory. However, one could similarly write the GW_{4D} model as a group field theory on $U(1)^4$. From this perspective, it seems interesting to investigate in the future the relationship between non-commutative field theories of the Moyal-type and group field theories. A possible approach could be to push further the relation between the Moyal star-product and the non-commutativity based on group manifolds as investigated in [19].

ACKNOWLEDGEMENTS

Discussions with Razvan Gurau are gratefully acknowledged. Research at Perimeter Institute is supported by the Government of Canada through Industry Canada and by the Province of Ontario through the Ministry of Research and Innovation.

APPENDIX: ONE-LOOP β-FUNCTION OF THE $(p^{\frac{1}{2}+\epsilon} + q^{\frac{1}{2}})$–MODEL

We prove in this appendix that all the models with propagators of the form $p^{\frac{1}{2}+\epsilon} + q^{\frac{1}{2}}$, $\epsilon \in]0, 1/2]$ are asymptotically free in the UV. The calculation of the β-function is made for $\epsilon = 1/2$ corresponding to the $\sigma T3$ model. For the remaining models, the proof is totally similar.

The β-function is encoded in the ratio

$$\lambda^{\text{ren}} = -\frac{\Gamma_4(0,0,0,0)}{Z^2}\,, \tag{A.1}$$

where λ^{ren} is the renormalized coupling (and so λ stands for the bare coupling). $\Gamma_4(m,n,p,q)$ is the amputated four-point function truncated at one-loop and which should be computed at zero external momenta in (A.1). Z is the wave function renormalization which should involve the subleading log-divergent term obtained after the Taylor expansion of the self-energy Σ which is the sum of the amputated two-point functions truncated at one-loop.

Σ can be evaluated as in the ordinary situation [10] and involves two types of contributions called tadpoles "up" T^+ and "down" T^-(see Fig.4).

We have

$$\Sigma = A_{T^+} + A_{T^-} = \frac{(-\lambda)}{2}\Big[2\sum_k \frac{1}{m+\sqrt{k}+\mu} + 2\sum_k \frac{1}{\sqrt{n}+k+\mu}\Big]\,. \tag{A.2}$$



FIG. 4. Tadpoles up T^+ and down T^- and four-point graph F.

As noticed, $\Sigma = \Sigma(m, \sqrt{n})$, so that evaluating $\partial_m \Sigma$ or $\partial_{\sqrt{n}}\Sigma$, we only collect the log-divergent part contributing to the wave function renormalization and this is

$$Z = 1 - \partial_m \Sigma|_{m=0} = 1 - \lambda S\,, \qquad S = \sum_k \frac{1}{(\sqrt{k}+\mu)^2}\,. \tag{A.3}$$

To Γ_4 contribute only a unique divergent four-point function F of the form given by Fig.4. We have

$$\Gamma_4(m,n,p,q) = -\lambda + \frac{1}{2}\frac{\lambda^2}{2^2}(2\cdot 2\cdot 2)\sum_k \frac{1}{(m+\sqrt{k}+\mu)(p+\sqrt{k}+\mu)}\,. \tag{A.4}$$

Note that, at this level, the above model differs from the GW 4D model since more graphs contribute to the Γ_4 function in the latter inducing a combinatorial factor twice greater.

We are in position to compute the β-function:

$$\lambda^{\rm ren} = -\frac{\Gamma_4(0,0,0,0)}{Z^2} = -\frac{(-\lambda+\lambda^2 S)}{(1-\lambda S)^2} = \lambda + \lambda^2 S\,. \tag{A.5}$$

Therefore $\beta = +1$ and $\sigma T3$ is asymptotically free in the UV as expected from [2]. In a similar way, all theories with propagator $p^{\frac{1}{2}+\epsilon'} + q^{\frac{1}{2}}$, $\epsilon' \in]0, 1/2]$ will be asymptotically free (once again, by the same reasons, T^- should be dropped because $\sum_k 1/(k^{\frac{1}{2}+\epsilon}+\mu)^2$ is convergent and the same contributions of the four-point function are still convergent and should be neglected) whereas, at the end-point, $\epsilon' = 0$, the GW_{4D} model becomes safe.

---

[1] J. Ben Geloun and V. Rivasseau, "A Renormalizable 4-Dimensional Tensor Field Theory," arXiv:1111.4997 [hep-th].

[2] J. Ben Geloun and D. O. Samary, "3D Tensor Field Theory: Renormalization and One-loop β-functions," arXiv:1201.0176 [hep-th].

[3] J. Ben Geloun, "Two and four-loop β-functions of rank 4 renormalizable tensor field theories," arXiv:1205.5513 [hep-th].

[4] D. Oriti, "The group field theory approach to quantum gravity," arXiv:gr-qc/0607032.

[5] V. Rivasseau, "Towards Renormalizing Group Field Theory," PoS C **NCFG2010**, 004 (2010) [arXiv:1103.1900 [gr-qc]].

[6] V. Rivasseau, "Quantum Gravity and Renormalization: The Tensor Track," arXiv:1112.5104 [hep-th].

[7] V. Rivasseau, "From perturbative to constructive renormalization," Princeton series in physics (Princeton Univ. Pr., Princeton, 1991).

[8] R. Gurau, "The complete 1/N expansion of colored tensor models in arbitrary dimension," Annales Henri Poincare **13**, 399 (2012) [arXiv:1102.5759 [gr-qc]].

[9] V. Bonzom, R. Gurau, A. Riello and V. Rivasseau, "Critical behavior of colored tensor models in the large N limit," Nucl. Phys. B **853**, 174 (2011) [arXiv:1105.3122 [hep-th]].

[10] V. Rivasseau, "Non-commutative renormalization," arXiv:0705.0705 [hep-th].

[11] H. Grosse and R. Wulkenhaar, "Renormalisation of phi**4 theory on noncommutative R**4 in the matrix base," Commun. Math. Phys. **256**, 305 (2005) [arXiv:hep-th/0401128].

[12] H. Grosse and R. Wulkenhaar, "Renormalization of phi**4 theory on noncommutative R**2 in the matrix base," JHEP **0312**, 019 (2003) [hep-th/0307017].

[13] P. Di Francesco, P. H. Ginsparg and J. Zinn-Justin, "2-D Gravity and random matrices," Phys. Rept. **254**, 1 (1995) [arXiv:hep-th/9306153].

[14] J. Ambjorn, B. Durhuus and T. Jonsson, "Three-Dimensional Simplicial Quantum Gravity And Generalized Matrix Models," Mod. Phys. Lett. A **6**, 1133 (1991).

[15] R. Gurau, "Colored Group Field Theory," Commun. Math. Phys. **304**, 69 (2011) [arXiv:0907.2582 [hep-th]].

[16] R. Gurau and J. P. Ryan, "Colored Tensor Models - a review," SIGMA **8**, 020 (2012) [arXiv:1109.4812 [hep-th]].

[17] V. Bonzom, R. Gurau and V. Rivasseau, "The Ising Model on Random Lattices in Arbitrary Dimensions," Phys. Lett. B **711**, 88 (2012) [arXiv:1108.6269 [hep-th]].

[18] V. Bonzom, R. Gurau and V. Rivasseau, “Random tensor models in the large N limit: Uncoloring the colored tensor models,” Phys. Rev. D **85**, 084037 (2012) [arXiv:1202.3637 [hep-th]].

[19] M. Dupuis, F. Girelli and E. R. Livine, “Spinors and Voros star-product for Group Field Theory: First Contact,” arXiv:1107.5693 [gr-qc].